

Riemann Paper(1859) Is False

Chun-Xuan. Jiang
P. O. Box3924, Beijing 100854, China
Jiangchunxuan@vip.sohu.com

Abstract

From Gamma function Remann derived the zeta function with Gamma function and put forward Riemann hypothesis(RH) which is false. The Jiang function $J_n(\omega)$ can replace RH. Green-Tao and Wiles papers are false [9-12]. Friedlander-Iwaniec and Heath-Brown papers are too long to understand their correctness[14, 15].

Riemann defined the Riemann zeta function (RZF)[1]

$$\zeta(s) = \prod_P (1 - P^{-s})^{-1} = \sum_{n=1}^{\infty} \frac{1}{n^s}, \quad (1)$$

where $s = \sigma + ti, i = \sqrt{-1}$, σ and t are real, P ranges over all primes. RZF is the function of the complex variable s in $\sigma \geq 0, t \neq 0$, which is absolutely convergent.

In 1896 J. Hadamard and de la Vallee Poussin proved independently [2]

$$\zeta(1+ti) \neq 0. \quad (2)$$

In 1998 Jiang proved [3]

$$\zeta(s) \neq 0, \quad (3)$$

where $0 \leq \sigma \leq 1$.

Theorem 1. Riemann paper (1859) is false [1]

We define Gamma function

$$\Gamma\left(\frac{s}{2}\right) = \int_0^{\infty} e^{-t} t^{\frac{s}{2}-1} dt. \quad (4)$$

For $\sigma > 0$. On setting $t = n^2 \pi x$, we observe that

$$\pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) n^{-s} = \int_0^{\infty} x^{\frac{s}{2}-1} e^{-n^2 \pi x} dx. \quad (5)$$

Hence, with some care on exchanging summation and integration, for $\sigma > 1$,

$$\pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) \bar{\zeta}(s) = \int_0^{\infty} x^{\frac{s}{2}-1} \left(\sum_{n=1}^{\infty} e^{-n^2 \pi x} \right) dx$$

$$= \int_0^{\infty} x^{\frac{s}{2}-1} \left(\frac{\mathcal{G}(x)-1}{2} \right) dx, \quad (6)$$

where $\bar{\zeta}(s)$ is called Riemann zeta function with gamma function rather than $\zeta(s)$,

$$\mathcal{G}(x) := \sum_{n=-\infty}^{\infty} e^{-n^2 \pi x}, \quad (7)$$

is the Jacobi theta function. The functional equation for $\mathcal{G}(x)$ is

$$x^{\frac{1}{2}} \mathcal{G}(x) = \mathcal{G}(x^{-1}), \quad (8)$$

and is valid for $x > 0$.

Finally, using the functional equation of $\mathcal{G}(x)$, we obtain

$$\bar{\zeta}(s) = \frac{\pi^{\frac{s}{2}}}{\Gamma\left(\frac{s}{2}\right)} \left\{ \frac{1}{s(s-1)} + \int_1^{\infty} (x^{\frac{s}{2}-1} + x^{\frac{s}{2}-\frac{1}{2}}) \cdot \left(\frac{\mathcal{G}(x)-1}{2} \right) dx \right\}. \quad (9)$$

From (9) we obtain the functional equation

$$\pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) \bar{\zeta}(s) = \pi^{-\frac{1-s}{2}} \Gamma\left(\frac{1-s}{2}\right) \bar{\zeta}(1-s). \quad (10)$$

The function $\bar{\zeta}(s)$ satisfies the following

1. $\bar{\zeta}(s)$ has no zero for $\sigma > 1$;
2. The only pole of $\bar{\zeta}(s)$ is at $s = 1$; it has residue 1 and is simple;
3. $\bar{\zeta}(s)$ has trivial zeros at $s = -2, -4, \dots$ but $\zeta(s)$ has no zeros;
4. The nontrivial zeros lie inside the region $0 \leq \sigma \leq 1$ and are symmetric about both the vertical line $\sigma = \frac{1}{2}$.

The strip $0 \leq \sigma \leq 1$ is called the critical strip and the vertical line $\sigma = \frac{1}{2}$ is called the critical line.

Conjecture (The Riemann Hypothesis). All nontrivial zeros of $\bar{\zeta}(s)$ lie on the critical line $\sigma = \frac{1}{2}$, which is false. [3]

$\bar{\zeta}(s)$ and $\zeta(s)$ are the two different functions. It is false that $\bar{\zeta}(s)$ replaces $\zeta(s)$, Pati proved that not all complex zeros of $\bar{\zeta}(s)$ lie on the critical line: $\sigma = 1/2$ [4].

Schadeck pointed out that the falsity of RH implies the falsity of RH for finite fields [5, 6]. RH is not directly related to prime theory. Using RH mathematicians prove many prime theorems which is false. In 1994 Jiang discovered Jiang function $J_n(\omega)$ which can replace RH, Riemann zeta function and L-function in view of its proved feature: if $J_n(\omega) \neq 0$ then the prime equation has infinitely many prime solutions; and if $J_n(\omega) = 0$, then the prime equation has finitely many prime solutions. By using $J_n(\omega)$ Jiang proves about 600 prime theorems including the Goldbach's theorem, twin prime theorem and theorem on arithmetic progressions in primes[7,8].

In the same way we have a general formula involving $\bar{\zeta}(s)$

$$\begin{aligned} \int_0^\infty x^{s-1} \sum_{n=1}^\infty F(nx) dx &= \sum_{n=1}^\infty \int_0^\infty x^{s-1} F(nx) dx \\ &= \sum_{n=1}^\infty \frac{1}{n^s} \int_0^\infty y^{s-1} F(y) dy = \bar{\zeta}(s) \int_0^\infty y^{s-1} F(y) dy, \end{aligned} \quad (11)$$

where $F(y)$ is arbitrary.

From (11) we obtain many zeta functions $\bar{\zeta}(s)$ which are not directly related to the number theory.

The prime distributions are regular rather than random. The arithmetic progressions in primes are not directly related to ergodic theory ,harmonic analysis, discrete geometry, and combinatorics. Using the ergodic theory Green and Tao prove that there exist infinitely many arithmetic progressions of length k consisting only of primes which is false [9, 10, 11]. Fermat's last theorem is not directly related to elliptic curves. In 1994 using elliptic curves Wiles proved Fermat's last theorem which is false [12]. There are Pythagoras's theorem and Fermat's last theorem in the complex hyperbolic functions and complex trigonometric functions. In 1991 without using any number theory Jiang proved Fermat's last theorem which is Fermat's marvelous proof[7]. Using Jiang function $J_3(\omega)$ we prove the following theorems.

Theorem 2. Primes Represented by $P_1^n + mP_2^n$ [13]

(1) Let $n = 3$ and $m = 2$. We have

$$P_3 = P_1^3 + 2P_2^3.$$

We have Jiang function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3 - \chi(P)) \neq 0,$$

Where $\chi(P) = 2P - 1$ if $2^{\frac{P-1}{3}} \equiv 1 \pmod{P}$; $\chi(P) = -P + 2$ if $2^{\frac{P-1}{3}} \not\equiv 1 \pmod{P}$;

$\chi(P) = 1$ otherwise.

Since $J_n(\omega) \neq 0$, there exist infinitely many primes P_1 and P_2 such that P_3 is a prime.

We have the best asymptotic formula

$$\begin{aligned} \pi_2(N, 3) &= \left| \{P_1, P_2 : P_1, P_2 \leq N, P_1^3 + 2P_2^3 = P_3 \text{ prime}\} \right| \\ &\sim \frac{J_3(\omega)\omega}{6\Phi^3(\omega)} \frac{N^2}{\log^3 N} = \frac{1}{3} \prod_{3 \leq P} \frac{P(P^2 - 3P + 3 - \chi(P))}{(P-1)^3} \frac{N^2}{\log^3 N}. \end{aligned}$$

where $\omega = \prod_{2 \leq P} P$ is called primorial, $\Phi(\omega) = \prod_{2 \leq P} (P-1)$.

It is the simplest theorem. It is the Heath-Brown problem [14]. 84pages paper is too long to understand its correctness.

(2) Let $n = P_0$ be an odd prime, $2 \mid m$ and $m \neq \pm b^{P_0}$.

we have

$$P_3 = P_1^{P_0} + mP_2^{P_0}$$

We have

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3 - \chi(P)) \neq 0,$$

where $\chi(P) = -P + 2$ if $P \mid m$; $\chi(P) = (P_0 - 1)P - P_0 + 2$ if $m^{\frac{P-1}{P_0}} \equiv 1 \pmod{P}$;

$\chi(P) = -P + 2$ if $m^{\frac{P-1}{P_0}} \not\equiv 1 \pmod{P}$; $\chi(P) = 1$ otherwise.

Since $J_n(\omega) \neq 0$, there exist infinitely many primes P_1 and P_2 such that P_3 is a prime.

We have

$$\pi_2(N, 3) \sim \frac{J_3(\omega)\omega}{2P_0\Phi^3(\omega)} \frac{N^2}{\log^3 N}.$$

Theorem 3. The Polynomial $P_1^n + (P_2 + 1)^2$ Captures Its Primes [13]

(1) Let $n = 4$, We have

$$P_3 = P_1^4 + (P_2 + 1)^2,$$

We have Jiang function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3 - \chi(P)) \neq 0,$$

Where $\chi(P) = P$ if $P \equiv 1 \pmod{4}$; $\chi(P) = P - 4$ if $P \equiv 1 \pmod{8}$; $\chi(P) = -P + 2$ otherwise.

Since $J_n(\omega) \neq 0$, there exist infinitely many primes P_1 and P_2 such that P_3 is a prime.

We have the best asymptotic formula

$$\pi_2(N, 3) = \left| \{P_1, P_2 : P_1, P_2 \leq N, P_1^4 + (P_2 + 1)^2 = P_3 \text{ prime}\} \right|$$

$$\sim \frac{J_3(\omega)\omega}{8\Phi^3(\omega)} \frac{N^2}{\log^3 N}.$$

It is the simplest theorem. It is Friedlander-Iwaniec problem [15]. 95pages paper is too long to understand, why did Ann. of Math publish this paper? Because a mathematician featuredly reviewed that this paper is a monumental breakthrough.

(2) Let $n = 4m$, We have

$$P_3 = P_1^{4m} + (P_2 + 1)^2,$$

where $m = 1, 2, 3, \dots$.

We have Jiang function

$$J_3(\omega) = \prod_{3 \leq P \leq P_1} (P^2 - 3P + 3 - \chi(P)) \neq 0,$$

where $\chi(P) = P - 4m$ if $8m \mid (P - 1)$; $\chi(P) = P - 4$ if $8 \mid (P - 1)$; $\chi(P) = P$ if $4 \mid (P - 1)$;

$\chi(P) = -P + 2$ otherwise.

Since $J_3(\omega) \neq 0$, there exist infinitely many primes P_1 and P_2 such that P_3 is a prime. It is a generalization of Euler proof for the existence of infinitely many primes.

We have the best asymptotic formula

$$\pi_2(N, 3) \sim \frac{J_3(\omega)\omega}{8m\Phi^3(\omega)} \frac{N^2}{\log^3 N}.$$

(3) Let $n = 2b$, We have

$$P_3 = P_1^{2b} + (P_2 + 1)^2,$$

where b is an odd.

We have Jiang function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3 - \chi(P)) \neq 0,$$

Where $\chi(P) = P - 2b$ if $4b \mid (P - 1)$; $\chi(P) = P - 2$ if $4 \mid (P - 1)$; $\chi(P) = -P + 2$

otherwise.

We have the best asymptotic formula

$$\pi_2(N, 3) \sim \frac{J_3(\omega)\omega}{4b\Phi^3(\omega)} \frac{N^2}{\log^3 N}.$$

(4) Let $n = P_0$, We have

$$P_3 = P_1^{P_0} + (P_2 + 1)^2.$$

where P_0 is an odd. Prime.

We have Jiang function

$$J_3(\omega) = \prod_{3 \leq P} (P^2 - 3P + 3 - \chi(P)) \neq 0,$$

Where $\chi(P) = P_0 + 1$ if $P_0 | (P - 1)$; $\chi(P) = 0$ otherwise.

Since $J_3(\omega) \neq 0$, there exist infinitely many primes P_1 and P_2 such that P_3 is also a prime.

We have the best asymptotic formula

$$\pi_2(N, 3) \sim \frac{J_3(\omega)\omega}{2P_0\Phi^3(\omega)} \frac{N^2}{\log^3 N}.$$

The Jiang function $J_n(\omega)$ is closely related to the prime distribution. Using $J_n(\omega)$ we are able to tackle almost all prime problems in the prime distributions.

References

- [1] B. Riemann, *Über die Anzahl der Primzahlen unter einer gegebenen Grösse*, Monatsber Akad. Berlin, 671-680 (1859).
- [2] P. Borwein, S. Choi, B. Rooney, *The Riemann hypothesis*, pp28-30, Springer-Verlag, 2007.
- [3] Chun-Xuan. Jiang, *Disproofs of Riemann hypothesis*, Algebras Groups and Geometries 22, 123-136(2005). http://www.i-b-r.org/docs/Jiang_Riemann.pdf
- [4] Tribikram Pati, *The Riemann hypothesis*, arxiv: math/0703367v2, 19 Mar. 2007.
- [5] Laurent Schadeck, *Private communication*. Nov. 5. 2007.
- [6] Laurent Schadeck, *Remarques sur quelques tentatives de démonstration Originales de l'Hypothèse de Riemann et sur la possibilité De les prolonger vers une théorie des nombres premiers consistante*, unpublished, 2007.
- [7] Chun-Xuan. Jiang, *Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture*, Inter. Acad. Press, 2002. MR2004c: 11001, <http://www.i-b-r.org/Jiang.pdf>
- [8] Chun-xuan. Jiang, *The simplest proofs of both arbitrarily long arithmetic progressions of primes*, Preprint, Oct. 2006.
- [9] B. Kra, *The Green-Tao theorem on arithmetic progressions in the primes: an ergodic point of view*, Bull. Am. Math. Soc. 43, 3-23(2006).
- [10] B. Green and T. Tao, *The primes contain arbitrarily long arithmetic progressions*. To appear, Ann. of Math.
- [11] T. Tao, *The dichotomy between structure and randomness, arithmetic progressions, and the primes*. In proceedings of the international congress of mathematicians (Madrid. 2006). Europ. Math. Soc. Vol.1, 581-609, 2007.
- [12] A. Wiles, *Modular elliptic curves and Fermat's last theorem*, Ann. of Math. 141, 443-551

(1995)

- [13] Chun-Xuan. Jiang, Prime theorem in Santilli's isonumber theory (II), Algebras Groups and Geometries 20, 149-170(2003).
- [14] D. R. Heath-Brown, Primes represented by $x^3 + 2y^3$. Acta Math. 186, 1-84(2001).
- [15] J. Friedlander and H. Iwaniec, The polynomial $x^2 + y^4$ captures its primes. Ann. of Math. 148, 945-1040(1998).